

صحة بيئية وبيئة

e

# CONTENTS

# WEJ

Journal of the Chartered Institution of Water and Environmental Management



Volume 19 September 2005 No. 3

## PEER REVIEWED PAPERS

- 154** SEASONAL VARIATION OF BACTERIOLOGICAL AND CHEMICAL QUALITY OF DRINKING WATER: A CASE STUDY IN A PALESTINIAN DISTRICT  
*I. A. Al-Khabit, F. Daoud, F. Rasmawi, S. Wa'ra, and M. Kassabry*
- 159** IMPROVING THE PERFORMANCE OF GRANULAR ACTIVATED CARBON (GAC) VIA PRE-REGENERATION ACID TREATMENT  
*L. Perry, D. Essex, P. Giess, N. Graham, K. Kaur, S. Lambert and C. Spencer*
- 167** ROBERTSBRIDGE FLOOD ALLEVIATION SCHEME: THE CHALLENGES OF A FAST-TRACK APPROACH  
*N. Baker, J. Palmer and G. Elswood*
- 174** AN INVESTIGATION INTO THE APPLICATION OF FLOATING REED BED AND BARLEY STRAW TECHNIQUES FOR THE REMEDIATION OF EUTROPHIC WATERS  
*P. Garbett*
- 181** PUBLIC PARTICIPATION - DRAWING THE BOUNDARIES  
*I. G. Bush, A. Gillson, M. Hamilton and M. Perrin*
- 189** MEMBRANE BIOREACTORS AND CONSTRUCTED WETLANDS FOR TREATMENT OF RENDERING PLANT WASTEWATER  
*C. A. O'Brien, M. Scholz, and G. L. McConnachie*
- 199** SOLVING AN URBAN RIVER EROSION PROBLEM ON THE TILMORE BROOK, HAMPSHIRE (UK)  
*A. Brookes, A. Chalmers, and R. Vivash*
- 207** AFFORDABILITY OF HOUSEHOLD WATER SERVICES IN GREAT BRITAIN  
*J. W. Sawkins and V. A. Dickie*
- 214** THE EFFECTS OF IMPROVING SEWAGE SLUDGE DIGESTION  
*W. P. Barber*
- 225** POLLUTANT BUILD UP & RUN OFF ON HIGHWAYS; EXPANDING THE CURRENT METHODOLOGY FOR ADDITIONAL DETERMINANDS - TECHNICAL NOTE  
*J. Patel and O. Drieu*
- 226** MILESTONES IN THE DEVELOPMENT OF MUNICIPAL WATER TREATMENT SCIENCE AND TECHNOLOGY IN THE 19TH AND EARLY 20TH CENTURIES: PART I  
*P. J. Purcell*
- 238** THE 1999 CRYPTOSPORIDIUM RISK ASSESSMENT EXERCISE IN ENGLAND AND WALES - A GROUNDWATER OVERVIEW  
*B. Morris and J. Cunningham*
- 248** OPERATIONAL EXPERIENCE FROM A SMALL FOOTPRINT LAMELLA AND BAFF PLANT IN ABERDEEN.  
*M. Jolly*
- 256** THE ROLE OF PRICING ON INTEGRATED WATER MANAGEMENT AT THE INDUSTRIAL PARK LEVEL: A CASE OF TEDA  
*Y. Geng*
- 264** ADVANCED LEACHATE TREATMENT AT THE STEWARTBY LANDFILL SITE  
*R. Toddington, M. Pankhania and E. C. Clark*

## YOUNG AUTHOR'S WINNING PAPER

- 272** QUANTIFYING THE UNQUANTIFIABLE: ECOHYDROLOGY ON THE OUSE WASHES  
*E. Simons*

## TOPICAL NOTES

- 280** INDUCTION OF NEW CIWEM PRESIDENT FOR 2005-2006
- 282** PRESIDENT'S AWARD 2005
- 283** ANNUAL GENERAL MEETING

<p>The Chartered Institution of Water and Environmental Management 15 John Street, London WC1N 2EB England Tel: 0207 831 3110 Fax: 0207 405 4967 Email: admin@ciwem.org Visit: www.ciwem.org</p> <p>Registered charity number: 1043409 ISSN: 0951-7359</p> <p>Executive director: Nick Reeves Director of policy: Justin Tobetham Policy officer: Harriet Greene Tel: 0207 831 3110 Email: h.greene@ciwem.org Editor in chief: Erika Yoraw Tel: 0870 3000 695 Email: wej.editor@leadmedia.co.uk Assistant editor: Wendy Hiles Tel: 0870 3000 690 Email: wendy@leadmedia.co.uk</p>	<p>Editorial panel: Dr Colin Powlesland (chairman), Charles Ruxton, Dr David Cook, Dr James Wishart, Dr Andrew Brookes, John Bridgeman, Mike Little, Dr Nicholas Robins, Paul Smith, Peter Dumble, Dr Peter Newman, Ross Gregory</p> <p>Published: In March, June, September &amp; December</p> <p>Annual subscriptions to non-members (UK) Water &amp; Environment Manager Magazine (WEM) &amp; Journal of the Chartered Institution of Water and Environmental Management (WEJ) ..... £248 Journal (WEJ) only ..... £165 Journal (WEJ) single issue UK ..... £42 CD ROM version of Journal (WEJ) ..... £155 Magazine (WEM) only ..... £98</p> <p>For further information about subscriptions Visit: www.ciwempublications.co.uk</p>	<p>Publisher: Gavin Touben Published by: Leadmedia Ltd 8 Harford Court, John Tate Road, Hertford Herts SG13 7NW Tel: 0870 3000 690 Fax: 0870 3000 691 Email: wej@leadmedia.co.uk</p> <p>Advertisement sales: Andrew Robinson, Leadmedia Ltd Tel: 0870 3000 690 Email: andrew@leadmedia.co.uk For further information about advertising Visit: www.ciwempublications.co.uk</p> <p>Printed on recycled paper V. Green, Regard paper 100% Post-consumer waste OF Chlorine free ISO 14001, ISO 9001 Produced at: Papeteries Mulsières &amp; Forest Mill, France © IPI/Priza of Innovation 2002 and 2003</p>
---	---	---

The Chartered Institution of Water and Environmental Management. The Institution is not responsible as a body for the statements made or opinions advanced in any of the papers or discussions published in its proceedings. Storage or reproduction of the contents, either as a whole, or in part, is forbidden unless specific permission has been obtained from the Institution. Reference to papers, or quotations there from, may be published elsewhere without permission, provided the exact source is acknowledged.



# SEASONAL VARIATION OF BACTERIOLOGICAL AND CHEMICAL QUALITY OF DRINKING WATER: A CASE STUDY IN A PALESTINIAN DISTRICT

I. A. Al-Khatib, F. Daoud, F. Rasmawi, S. Wa'rra AND M. Kassabry\*

## ABSTRACT

Evaluation of drinking water quality for Jenin district in Palestine was carried out for the years 1997, 1998, and 1999, respectively. Only 53.1%, 85.4% and 69.6% of the drinking water samples tested for free chlorine residual, total coliforms and faecal coliforms, respectively are within the limits of the Palestinian and International standards. The seasonal variation of bacteriological and chemical quality of drinking water was investigated. It was found out that the summer season has the best quality and the winter has the worst. There remains some ignorance in testing the drinking water in the villages and towns in Jenin district. Huge efforts are required to improve the drinking water quality in Jenin district, as well as other Palestinian districts, through public awareness, training of governmental inspectors, in addition to a strict monitoring system for water quality.

**Key words:** faecal coliforms; total coliforms; free chlorine residual; Jenin district; and Palestine.

\*Institute of Community and Public Health, Birzeit University, West Bank, Palestine

## INTRODUCTION

Water is an indispensable commodity, which should be easily accessible, adequate in quantity, free of contamination, safe, inexpensive and readily available throughout the year in order to sustain life. According to the World Health Organization (WHO<sup>(1)</sup>), and others<sup>(2,3,4,5)</sup>, protection of water supplies from contamination is the first line of defence.

According to the Palestine Standards Institution, PSI<sup>(6)</sup>, in order to have acceptable drinking water, we have to examine the water continuously to make sure that it is free from bacteria and should have the concentrations of substances such as free chlorine residual based on the guideline value. One must distinguish clearly between the examination of chlorinated and unchlorinated water supplies. If one is examining a chlorinated water supply one knows that, if the chlorination process is working correctly, all coliform organisms will have been killed<sup>(7,8)</sup>.

Most of the time, the source water usually meets hygiene standards. However, free chlorine residual concentration rapidly declines in the distribution system, as there is little protection against secondary microbiological contamination

of water in pipelines, and rechlorination is not practiced. This situation results in very low free chlorine residual at taps, and therefore gastrointestinal morbidity might appear<sup>(9)</sup>.

The majority of the population in the Jenin district construct their own cisterns to collect rain water from the roofs of their houses during the winter seasons and utilize it for their domestic consumption. Approximately 60% of the wastewater generated from households in the Jenin City is collected by sewer networks. Unfortunately, the cesspits or cesspools are the main methods of wastewater disposal in the Jenin district. Approximately 13% of the population in the Jenin district is connected with the sewage network. Another serious problem in the Jenin district is that water pipelines are laid adjacent to sewage lines causing, in many cases, wastewater infiltration to the drinking water<sup>(10)</sup>. According to Al-Kahah<sup>(11)</sup> and Othman<sup>(12)</sup> there are 76 communities in Jenin district, out of which only 34 communities are connected to public water networks.

This study was performed to ascertain the bacteriological quality of the drinking water in Jenin district, in Palestine, in addition to some of its chemical characteristics. The aim of this

research district is hazard to and faecal water in include team of tested in years of

## MATERIALS

During J water sa houses, r towns, a Bank of Environm of Heat hundred coliforms nine sam ensure th were tak but no quantific microbic availabl faecal c Number Lactose

The d analyzer Package some litr found in labelled the sam

## RESULTS

The val are show chlorine concen samples ppm. A drinking water, th 0.2 ppm consum availabl limits of



and 1999,  
coliforms  
seasonal  
summer  
king water  
nin district,  
dition to a

practiced. This  
at taps, and  
car.  
strict construct  
e roofs of their  
ze it for their  
re wastewater  
s collected by  
cesspools are  
e Jenin district.  
Jenin district is  
other serious  
elines are laid  
es, wastewater  
to Al-Qahah<sup>(11)</sup>  
nin district, out  
o public water  
acteriological  
n Palestine, in  
The aim of this

research is to determine if the drinking water throughout Jenin district is suitable for drinking and does not constitute a health hazard to the population. In this study, the distribution of total and faecal coliforms and free chlorine residual in the drinking water in Jenin district will be examined and analysed. This will include all the water samples which were collected by the team of environmental health inspectors from MoH and tested in the public health laboratory of the MoH during the years of 1997-1999.

**MATERIALS AND METHODS**

During January 1997 until 26 December 1999, 1964 drinking water samples were collected from different locations (wells, houses, restaurants, springs, schools etc.) from the villages, towns, a city, and a refugee camp in Jenin district in the West Bank of Palestine. The samples were collected by the Environmental Health Department of the Palestinian Ministry of Health for inspection purposes. One thousand nine hundred and thirteen samples were examined for total coliforms (247 for faecal coliforms). One thousand and sixty-nine samples were examined for free chlorine residual, to ensure the chemical quality of the drinking water. The samples were taken during the day time, normally in the early morning, but no specific time was indicated for this purpose. A quantitative descriptive design, describing the microbiological and chemical quality of drinking water available in records, has been used. Total coliforms and faecal coliforms were enumerated by the Most Probable Number, MPN, method inoculating series of five tubes, using Lactose Broth- Brilliant Green Bile Broth respectively<sup>(12)</sup>.

The data was coded and entered into the computer and analyzed using the statistical analytical system SPSS (Statistical Package for Social Sciences). It was found that there were some limitations for this research, as many missing items were found in the sampling form results, many samples were not labelled well and the location was not identified, and not all the samples were tested for free chlorine residual.

**RESULTS**

The values of all microbiological and chemical parameters are shown in Table 1. Out of the tested 1069 samples for free chlorine residual, 488 (45.7%) of the samples have a concentration less than 0.2 ppm (0-0.1), 568 (53.1%) of the samples between 0.2-0.8 ppm and 13 samples more than 0.8 ppm. According to the WHO<sup>(9)</sup> and PSI<sup>(7)</sup> guidelines for drinking water quality, for effective disinfection of drinking water, the concentration of free chlorine residual should be 0.2 ppm and not more than 0.8 ppm when water reaches the consumer. This means that only 53.1% of the drinking water available in Jenin District is acceptable for drinking within the limits of standards. This result directly reflects the presence of

microbiological contamination of the drinking water.

According to the WHO<sup>(9)</sup> and Al-Qahah<sup>(11)</sup> (2001) classification of total coliforms, out of the 1913, 85% of samples are free of contamination (zero degree of contamination), while 7.8% and 6.8% of samples are in the second and third degrees of contamination respectively. None of the examined water samples were within the third degree of contamination. Out of 247 samples tested for faecal coliforms, 172 (69.6%) were of good quality and had no risk, while 21 (8.5%) of the samples had low risk, 33 (13.4%) of the samples had an intermediate risk level, 21 (8.5%) of the water samples showed high risk level, and none of these samples showed very high risk level. This classification of risk is recommended by WHO<sup>(9)</sup>.

Table 1. Range Values, Frequencies and Valid Percentages of Free Chlorine Residual (FCR), Total Coliforms and Faecal Coliforms (TC-FC)

Range Values, Frequencies and Percentages								
Free Chlorine Residual (ppm)			TC (n/100ml)			FC (n/100ml)		
0-0.1	488	45.7	0-3	1633	85.4	0	172	69.6
0.2-0.8	568	53.1	4-50	150	7.8	1-10	21	8.5
>0.8	13	1.2	51-50,000	130	6.8	11-100	33	13.4
			>50,000	-	-	101-1000	21	8.5
						>1000	-	-
Total	1069	100.0		1913	100.0		247	100.0

Table 2 shows the results of sampling location versus annual distribution of population<sup>(10)</sup>, sample size, and sampling density per 1000 of population cross tabulations. The total number of samples shown in this table are 1570 out of 1964 tested, due to the limitations of the study. Nine hundred and fifty-two (60.6%) of the total samples were collected from the city of Jenin during the three years of 1997, 1998, and 1999; while 124 (7.9%) were from Jenin Refugee Camp. Four hundred and ninety-four (31.5%) of the samples were collected from the different villages and towns of Jenin District. The range of annual sampling densities in the City of Jenin, Jenin Refugee Camp and the villages and towns of Jenin district are between 10.3 -13.0, 3.9-5.1 and 2.4-3.0 samples per 1000 population respectively.

**DISCUSSION**

In order to see the effect of the concentration of free chlorine residual on the biological indicators, many cross tabulations have been performed. The final summary is presented in Table 3. As it is clearly seen from this table, there is an inverse relationship between presence of TC and FC and the concentration of free chlorine residual. Most of the contaminated samples with faecal coliforms have a free chlorine residual less than 0.2 ppm. Only one sample tested for

Table 2. Sampling Location Versus Annual Distribution of Population, Sample Size (Pop, N), and Sampling Density (1000 N/ Pop)

Sampling Location	Sampling Year								
	1997			1998			1999		
	Pop*	N	(1000 N/ Pop)	Pop*	N	(1000 N/ Pop)	Pop*	N	(1000N/ Pop)
Jenin City	26332	272	10.3	27329	354	13.0	28394	326	11.5
Jenin Refugee Camp	8991	35	3.9	9331	48	5.1	9695	41	4.2
Villages And Towns	157420	165	1.0	163381	203	1.2	169748	126	0.7
Total	192743	472	2.4	200041	605	3.0	207837	493	2.4

\*PCPS<sup>(14)</sup>

Table 3. Effect of Free Chlorine Residual Concentration (FCR) on Total Coliforms and Faecal Coliforms (TC, FC) and the Degree of Contamination Risk.

Free Chlorine Residual (ppm)		TC (n/100ml)			FC (n/100ml)			
		0-3 no risk	4-50 low risk	51-50,000 intermediate risk	0 no risk	1-10 low risk	11-100 intermediate risk	101-1000 high risk
		0-0.1	453	25	9	23	3	1
0.2-0.8	525	23	20	21	1	1	1	
> 0.8	12	1	-	6	-	-	-	
Total		990	49	29	50	4	2	1

faecal coliform and has a high risk effect. Forty-nine samples tested for TC and two samples tested for FC are found to be within the degree of intermediate risk.

Table 4 summarizes the sampling location versus range values and frequencies of FCR, TC, FC, respectively. It is noticed that most of the samples are tested in Jenin city even though its population consists of only 13.7% of the total population of Jenin District. There is some attention given to drinking water sampling in Jenin Refugee Camp, but most of the samples tested for FCR have a concentration less than 0.2 ppm, which is the minimum concentration recommended by PSI<sup>(6)</sup> and WHO<sup>(9)</sup>. It is clearly seen that there is some ignorance in testing the drinking water in the villages and towns in Jenin district even though their population consists of 81.7% of the total population, and most of the contaminated samples with TC are in the villages and towns.

From the cross tabulations as shown in Table 5, it was

obvious that there is a significant statistical relationship between the sampling year and the contamination of drinking water with FC (Chi square = 19.834, p= 0.001) and the free chlorine residual concentration (Chi square = 44.967, p= 0.000) respectively. Table 5 shows that in 1997 and 1998, 83.1% of the samples tested for FC had no risk. This situation was improved in 1999 where 90.5% of the samples had no risk of contamination. In 1997, 47.4% of the samples tested for FCR concentration were within the limits of standards (0.2-0.8 ppm). The situation improved in 1998, as 64.5% of the tested samples were found to have FCR concentration within the limits of standards, but this trend had changed in 1999 with only 43.9% of the samples within the limits of standards.

Regarding the FC, there is no significant relationship between the sampling year and degree of contamination, and this is clear in Table 5. For example, in 1997, 70.5% of the samples tested for FC showed no risk of contamination, while

Table 4. Sampling Location Versus Range Values and Frequencies of Free Chlorine Residual (FCR), Faecal Coliforms and Total Coliforms (TC, FC).

Sampling Location	Range values and frequencies									
	FCR (ppm)			TC (n/100ml)			FC (n/100ml)			
	0.0-0.1	0.2-0.8	>0.8	0-3	4-50	51-50,000	0	1-10	11-100	101-1000
Jenin City	286	355	7	875	41	34	18	-	-	-
Jenin Refugee Camp	94	7	1	115	7	2	1	1	-	-
Villages and Towns	34	157	-	432	47	12	9	1	2	-
Total	414	519	8	1422	95	48	28	2	2	-

Note: the numbers in the body of the table are the number of samples

Table 5. Cro anc

Sampling Year

1997

1998

1999

Total

Table 6. Cro Coli

Sampling Season

Summer

Autumn

Winter

Spring

Total

in 1998 and showed no the quality

In order quality, ma summary k table, the since 31.9% PSI<sup>(6-10)</sup> and was in the

within the li that the th of bacteric The sample risk, consists only 23.7% of risk. The there is a drinking investigation



Table 5. Cross Tabulations of Sampling Year versus Range Values, Frequencies and Valid Percentages of Free Chlorine Residual (FCR), Total Coliforms and Faecal Coliforms (TC -FC).

Sampling Year	Range Values, Frequencies and Percentages											
	Free Chlorine Residual (ppm)			TC (n/100ml)				FC (n/100ml)				
	0-0.1	0.2-0.8	>0.8	0-3	4-50	51-50,000	>50,000	0	1-10	11-100	101-1000	>1000
1997	119	108	1	507	50	53	-	67	8	9	11	-
	52.2	47.4	0.4	83.1	8.2	8.7		70.5	8.4	9.5	11.6	
1998	149	285	8	599	69	53	-	54	7	15	2	-
	33.7	64.5	1.8	83.1	9.6	7.4		69.2	9.0	19.2	2.6	
1999	220	175	4	527	31	24	-	51	6	9	8	-
	55.1	43.9	1.0	90.5	5.3	4.1		68.9	8.1	12.2	10.8	
Total	448	568	13	1633	150	130	-	172	21	33	21	-
	45.7	53.1	1.2	85.4	7.8	6.8		69.6	8.5	13.4	8.5	

Table 6. Cross Tabulations of Sampling Season versus Range Values, Frequencies and Valid Percentages of Free Chlorine Residual (FCR), Total Coliforms and Faecal Coliforms (TC -FC).

Sampling Season	Range Values, Frequencies and Percentages											
	Free Chlorine residual (ppm)			TC (n/100ml)				FC (n/100ml)				
	0-0.1	0.2-0.8	>0.8	0-3	4-50	51-50,000	>50,000	0	1-10	11-100	101-1000	>1000
Summer	155	181	9	442	47	44	-	63	5	14	4	-
	31.8	31.9	69.2	27.1	31.3	33.8		36.6	23.8	42.4	19.0	
Autumn	108	140	-	374	29	29	-	36	3	6	3	-
	22.1	24.6		22.9	19.3	22.3		20.9	14.3	18.2	14.3	
Winter	129	105	-	388	43	25	-	9	4	3	3	-
	26.4	18.5		23.7	28.7	19.2		5.2	19.0	9.1	14.3	
Spring	96	142	4	429	31	32	-	64	9	10	11	-
	19.7	25.0	30.8	26.3	20.7	24.6		37.2	42.9	30.3	52.4	
Total	488	568	13	1633	150	130	-	172	21	33	21	-
	100.0	100.0	100.0	100.0	100.0	100.0		100.0	100.0	100.0	100.0	

In 1998 and 1999, 69.2% and 68.9% of the samples tested for FC showed no risk of contamination, respectively. This means that the quality of drinking water has not improved over the years.

In order to see the seasonal variation of sampling and water quality, many cross tabulations have been performed. The final summary is presented in Table 6. As is clearly seen from this table, the summer season is the best for FCR concentration since 31.9% of the samples that are within the limitation of the PSI<sup>(6-15)</sup> and WHO<sup>(6)</sup> standards are in this season. The worst quality was in the winter season, since only 23.7% of the samples are within the limitation of the standards. It can be concluded also that the the summer season is the best from the point of view of bacteriological quality of the tested samples (see Table 6). The samples tested for TC in the summer, with zero degree of risk, consists of 27.1% of the 1633 samples tested for TC. In winter only 23.7% of the 1633 samples tested for TC are of zero degree of risk. The same result can be concluded for FC. That means there is a strong relationship between the disinfection of drinking water and its bacteriological quality. More investigation is required to know the reasons for the bad quality

of drinking water mainly in winter.

It was noticed that none of the samples were examined for faecal streptococci or sulphite-reducing clostridia, as recommended by PSI<sup>(6)</sup> when samples are contaminated with faecal coliform. According to WHO<sup>(6)</sup> and PSI<sup>(6)</sup> it is recommended that the drinking water should not contain enteric viruses, three of the intestinal protozoa that are pathogenic for man and can be transmitted by drinking water: Entamoeba histolytica, Giardia Spp. and Balantidium coli - all helminthes that can be infective to man and free-living organisms, including plankton and macro invertebrates. It was noticed that none of the samples were examined for any of the above mentioned indicators.

## CONCLUSIONS

From the analysis of the results the following conclusions can be drawn:

1. The Ministry of Health in Jenin district focussed attention on the city more than other areas in the same district, such as the camp, the villages and the towns.

2. The range of annual sampling densities in the City of Jenin, Jenin Refugee Camp and the villages and towns of Jenin District, are between 10.3 -13.0, 3.9-5.1 and 2.4-3.0 samples per 1000 population, respectively.

3. There is a significant statistical relationship between the sampling year and the contamination of drinking water with FC and the free chlorine residual concentration.

4. The seasonal variation of sampling indicated that the summer season is the best for water quality and the worst quality was in the winter season.

5. There is a strong relationship between the disinfection of drinking water and its bacteriological quality.

The following recommendation can be made as a result of the previous analysis:

1. More attention should be paid for sampling of drinking water in the villages and towns in Jenin district.

2. Adequate monitoring network should be established.

3. A time schedule for the biological and chemical examination of drinking water should be followed.

4. Public awareness about safety of drinking water and its impact on health should be emphasized.

5. Intersectoral collaborations between the Ministry of Health, the Ministry of Education and other related leaders should be established for planning a national strategy for providing safety water and improving the quality of water in Palestine.

## REFERENCES:

- (1) World Health Organization, WHO, *Guidelines for Drinking Water Quality: Health Criteria and Other Supporting Information*, Geneva, Vol 2, 1986.
- (2) MCFETERS, G. A., AND TORANZOS, G. A., *Detection of Indicator Microorganisms in Environmental Freshwaters and Drinking Water*. In: HURST, C. J. KNUDSEN, G. R. MCLNERNEY, M. J., STETZENBACH L. D AND WALTER M. V. (eds), *Manual of Environmental Microbiology*, 1997, U.S.A.
- (3) World Health Organization, WHO, *Guidelines for Drinking Water Quality: Surveillance and Control of Community Supply*, 2nd edition, Vol. 2, Geneva, 1997.
- (4) World Health Organization, WHO, *Guidelines for Drinking Water Quality: Recommendations*, 2nd edition, Vol. 1, Geneva, 1993.
- (5) MOE, C. L. Water borne Transmission of Infectious Agents. In: HURST, C. J. KNUDSEN, G. R. MCLNERNEY, M. J., STETZENBACH L. D AND WALTER M. V. (eds), *Manual of Environmental Microbiology*, U.S.A. 1997.
- (6) Palestinian Standards Institution, PSI, *Drinking Water*, PS41, Nablus, Palestine, 1997.
- (7) World Health Organization, WHO, *International Standards for Drinking Water*, 3rd edition, Geneva, 1971
- (8) World Health Organization, WHO, *Guidelines for Drinking Water Quality, Health Criteria and Other Supporting Information*, Vol. 2, Geneva, 1984.
- (9) ANDREY E., TIMOTHY F., ANDREY T., NINA D., IRENA S. AND VALDISLAV F., Deterioration of drinking water quality in the distribution system and gastrointestinal morbidity in a russian city, *Int. J. of Env. Health Res.*, 12 (3), **221-233**, U.K, 2002.
- (10) Applied Research Institute-Jerusalem, ARIJ, *Environmental Profile for the West Bank Volume 7, Jenin District, Bethlehem, Palestine*, 1996.
- (11) AL-KAHAH, A., *Water-Borne Pathogens with Relation to Gastroenteritis in Salfeet District: an-Epidemiological Study*, MSc. thesis, An-Najah National University, Nablus, Palestine, 2001.
- (12) OTHMAN, S., *Bacterial Quality of Drinking Water in Rain-Fed Cisterns and Roof Storage Tanks in Beit Lied and Safarine Villages*, MSc. thesis, An-Najah National University, Nablus, Palestine, 2000.
- (13) American Public Health Association, APHA, American Water Works Association, Water Environment Federation, *Standard Methods for the Examination of Water and Wastewater*, 18th ed. Washington, DC, U.S.A., 1992.
- (14) Palestinian Central Bureau of Statistics, PCBS, *Small Area Population*, 1997-2010, Ramallah, Palestine, 1999.
- (15) AL- KHATIB I. A., KAMAL S., TAHA B., AL HAMAD J., AND JABER H., Water- health relationships in developing countries: a case study in Tulkarem district in Palestine, *International Journal of Environmental Health Science*, U. K., 13, **199-206**, 2003.

GI

AB

A la  
calc  
rege  
rem  
trea  
surc  
was  
whic  
that

Key

\*Wa

This pa

INTRC

The intrc  
Water St  
control  
research  
water tr  
carbon  
and ad: